

MOLTRAQ 芸 š ()

MOLTRAQ

Evolution of VHS in Denmark Susie Sommer Mikkelsen Annual Workshop 2015





DTU Vet National Veterinary Institute

VHS disease description

- Notifiable disease in Europe
- \bullet Typically occurs spring/autumn at low temperatures (10-14 $^{\circ}\,$ C)
- Only been found in the Northern hemisphere

DTU



VHS symptoms

- Signs of infection include hemorrhaging of skin, muscles and internal organs
- Other symptoms: Pale gills, exophthalmos and listless behaviour
- Est. Mortality: 80-100% in fry and fingerling 50% in large adults





Susceptible species



Danish Fish farm description



■ 5 - 100 tonnes ■ > 100 tonnes



No. Of fishfarms according to size (2013):

< 5 tonnes	5 - 100 tonnes	> 100 tonnes	all in all	
19	84	140	243	

VHS Denmark history



- · First discovered in 1963
- Economical losses of up to nearly 40 mill. Euro/year
- Eradication progamme started in 1965
- Mainly conducted by emptying and draining fish farms removing all fish and water
- This was followed by extensive cleaning and disinfection of the whole
- Restocking only with fish from approved disease free farms
- Electrofishing of wild fish from streams



VHS Denmark history


- No cases since early 2009
- Denmark officially free of VHS in 2013, but:
- Still VHSV in marine fish in the surrounding marine environment



VHS Virus description




- Belongs to the genus Novirhabdovirus and the Rhabdoviridae family
- · Closely related to IHNV
- Bullet-shaped virus



DTU

VHS Virus description

- Non-segmented negative-sense single-stranded RNA genome
- Genome is 11-12 kb in size
- Encodes five structural proteins and one non-structural protein



Dataset description




Time-period: 1962-2009

Epidemiological information includes:

- Sampling date
- Clinical status
 Geographical information incl.
 - Coordinates
 - · River systems
- Farms positioning relative to each other and to the top of the river
 Historical knowledge about the different outbreaks and farms



Distribution of isolates





Distribution of isolates


Dataset description - sequences





- Full-length G-gene sequenced
- 309 isolates in dataset
- 221 isolates sequenced in the MOLTRAQ project, 10 isolates not sequenced
- 136 unique sequences
- 173 sequences with at least 1 identical sequence
- 0-4,6% difference between isolates
































Thank you

- Henrik Korsholm, Danish Veterinary and Food Administration
- My colleagues at DTU:
 Morten S. Bruun
 Niccoló Vendramin
 Torsten Boutrup
 Niels-Jørgen Olesen











25 DTU Vet, Technical University of Denmark

Frederiksberg 28/05-2015